

Krystexxa (pegloticase) | Order Form

Patient Name: _____ DOB: _____ Phone: _____
Address: _____ City: _____ State: _____ Zip: _____

1. For new patients, please submit with form:

- Copy of insurance card
- Patient demographics
- History & physical
- Labs/records including G6PD deficiency screening (if indicated*) and baseline uric acid levels

2. Patient Information

Male Female Height: _____ in/cm Weight: _____ lbs/kg Allergies: _____

Is this the first dose? Yes No, date of last infusion: _____ Line type: PIV PICC Port Other

3. Diagnosis and Clinical Information

Primary diagnosis information: Gout Other: _____ ICD-10 (required): _____ Is there an immunomodulator prescribed? Yes No **If yes, please indicate:** Methotrexate Other _____ Is patient currently taking oral urate-lowering agents? No Yes. Oral urate-lowering agents should be discontinued prior to Krystexxa

G6PD deficiency screening and/or testing results:

- Patient evaluated by provider and is **not** at risk and will **not** be tested (or tested negative)
- Patient is at risk for G6PD deficiency, and test results are: Positive (contraindicated) Negative

Baseline serum uric acid level: _____ mg/dL (Must be >6.0mg/dL to initiate Krystexxa)

- Uric acid levels are required to be drawn 1 to 2 days prior to each infusion, **referring provider to arrange lab draws locally**
Name of lab facility: _____ Phone number for lab results: _____
- Please fax results to Coastal Infusion Services at 985-792-9004 as soon as available
- A single uric acid of >6.0mg/dL will require follow-up with provider, but will not post-pone next infusion

4. Prescription Information

Medication	<input checked="" type="checkbox"/> Krystexxa (pegloticase)
Dosing / Frequency	<input checked="" type="checkbox"/> 8mg in 250mL sodium chloride 0.9% IV every 2 weeks
Administration	<input checked="" type="checkbox"/> Prepare and infuse per manufacturer guidelines. Infuse over no less than 2 hours and observe patient for at least 1 hour following infusion <input checked="" type="checkbox"/> May infuse in patient home unless otherwise noted: _____
Quantity / Refills	Dispense 2-week supply on all selected medications; Refill x 12 months unless otherwise specified: _____ Dispense all medical supplies necessary for infusion

5. Additional Orders

RN to start peripheral IV or use existing CVC. RN to administer catheter flushing per Coastal Infusion Services Policy and Procedure

Give standard premedications 30 minutes prior to infusion:

Solu-medrol 40 mg **OR** _____ mg IV

Acetaminophen 650mg PO

Antihistamine: patient may take night prior AND morning of infusion. If nothing is checked, patient/pharmacist may select from the following OTC products:

Diphenhydramine 25mg PO

Fexofenadine 60 mg
(Allegra 12 Hour) PO

Cetirizine 10mg PO

Fexofenadine 180 mg
(Allegra 24 Hour) PO

Loratadine 10mg PO

Other: _____

RN to instruct patient to hydrate pre/post infusion and educate on taking OTC diphenhydramine and/or acetaminophen per manufacturer dosing recommendations as needed to prevent/treat post-infusion headache.

RN to monitor patient for at least 1 hour post infusion and educate on possible side effects, allergic reactions, and when to contact physician

6. Adverse Reaction Orders

Standard anaphylaxis kit to be dispensed and dosed per protocol: Epinephrine IM/SQ (1 mg/mL vial), diphenhydramine IV/IM (50 mg/mL vial), and NS IV. Additional orders: _____

7. Prescriber Information

Prescriber Name: _____ Office Contact: _____

Address: _____ City: _____ State: _____ Zip: _____

Phone: _____ Fax: _____ License No.: _____

DEA No.: _____ NPI: _____

Physician Signature (Substitution Permitted)

Date

Physician Signature (Dispense as Written)

Date